

When does a broken heart become a mental disorder?

Rarely, if ever...

But don't tell that to the American Psychiatric Association, which has just released its fifth version of the Diagnostic and Statistical Manual of Mental Disorders. The DSM is a catalogue of mental health disorders, hundreds of them, each thralling a listing of symptoms. The manual informs selection of a diagnosis which is required by U.S. insurance companies for reimbursement for medical health care.

There's a major change in the newest version, DSM-5 with serious implications for the millions of people who are coping with the death of a child, spouse, parent, friend or other loved one.

But first, a quick glimpse of the history of this publication, often referred to as the bible of psychiatry. The very first edition published in 1952 didn't even refer to grief, considering it an accepted and normal reaction to the death of a loved one. The third edition added an exclusion statement under Major Depressive Disorder, referred to as the 'bereavement exclusion'. Under this exclusion a diagnosis of Major Depressive Disorder could not be made for a full year after a death. They recognised that normal and common reactions to the death of a loved one could look like symptoms of depressive disorder, for example sadness, disturbed sleep, lack of concentration changes in eating and loss of interest in things that were once pleasurable.

In 1994 the fourth version of the DSM reduced the bereavement exclusion to two months after a death and this new version removes the bereavement exclusion completely, meaning in effect that anyone can receive a diagnosis of Major Depressive Disorder two weeks after the death of a child, parent, spouse, friend or anyone.

Why does this matter?

For at least three reasons.

First, normal reactions to the death of a loved one will be easily misclassified as the mental disorder depression. Grief is not the same experience as a major depressive disorder. It is not an illness to be treated or cured. It is a healthy response to a painful reality that one's world is forever altered, and will never be the same. Absorbing this loss, and adapting to all the changes it unleashes, has its own unique course for every person, and will not be stilled or stopped by quick fixes or simple solutions. Death is a life-altering event, but grief is not a pathological condition.

Second, antidepressants are commonly and frequently prescribed. There is a strong likelihood that newly bereaved people will qualify for a diagnosis of Major Depressive Disorder just two weeks after a death even though their reactions are normal. Antidepressants have not been shown to be helpful with grief-related depressive symptoms, and there is accumulating evidence of long-term negative effects of being on antidepressants. We need to ask why psychiatry is pathologising grief and therefore making

inappropriate pharmacological treatment easier. And we should not overlook the self-interest of pharmaceutical companies who see a new and substantial market for antidepressants, currently a multi-billion dollar industry.

Third, about 80% of prescriptions for antidepressants are written by medical doctors, not psychiatrists. We have the expectation that physicians, as well as psychologists, social workers and clergy, to whom many of us turn for help after losses of all kinds, have professional training, solid research backing and supervised experience to guide them. Some do, but in fact, a considerable majority of practitioners with these degrees have no professional training at all in responding to the bereaved.

The caution here? Be wary of physicians or other medical professionals who rush to prescribe antidepressants to address your grief.

Here's a better prescription. Mourn the death of your loved one in your own way. There is no prescribed formula. You may cry, you may not. Your reactions will be shaped by many things; the relationship you had with the deceased, your personality style and the support or lack of support you receive from others. Push aside those who tell you to move on, that every cloud has a silver lining. What one person finds comforting may not work for another. Find friends and family who understand, and with whom you can share your experience. If they won't listen or help, or if their help is not enough, search for support groups through your local hospital, hospice or community organisations. Don't be afraid to seek professional help, but if you do, ask about the person's training, qualifications and experience with grief, loss and bereavement.

We grieve as deeply as we love. We can get off track with love and we can respond to our grief in ways that aren't healthy or don't serve us well. But let's not make love, or grief a mental disorder.

This document was written by a group of concerned professionals in response to the release of the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders (DSM-5)

Thomas Attig, PhD, Professor Emeritus in Philosophy, Bowling Green State University

Inge B. Corless, RN, PhD, FAAN, Professor, MGH Institute of Health Professionals Boston MA

Kathleen R. Gilbert, PhD, Executive Associate Dean, Indiana University School of Public Health, Bloomington, IN

Dale G. Larson, PhD, Professor, Department of Counseling Psychology, Santa Clara University, CA

Mal McKissock, OAM, Director of Clinical Services, Bereavement Care Centre, Sydney, Australia

David Roth, Executive Director, Puetz-Roth Funerals and Grief Companions, Bergisch Gladbach, Germany

Donna Schuurman, EdD, FT, Executive Director, The Dougy Centre for Grieving Children and Families, Portland, OR

Phyllis R. Silverman, PhD, Professor Emerita, Scholar-in-Residence, Women's Studies

Research Centre, Brandeis University, Waltham, MA
J. William Worden, PhD, ABPP, Psychologist, Laguna Niguel, CA

We would like to acknowledge the international Work Group on Death, Dying and Bereavement (IWG) for the opportunity to develop these ideas. This statement represents the opinions of the authors, not the opinions of the Board or membership of the IWG